

Webs & Smooth Components of Two Column Springer Fibers


Mike Cummings

Department of Combinatorics and Optimization
University of Waterloo

BIRS
Cluster Algebras, Webs, and Canonical Bases
25 March 2026

Springer's Resolution

Springer ('69) gave a resolution of singularities



of the nilpotent cone in \mathfrak{sl}_n

Spaltenstein ('76). The fiber of the resolution at η

$$\text{"Springer fiber"} \quad S_\eta = \{ \text{flags } V_\bullet \mid \eta \cdot V_i \subseteq V_{i-1} \text{ for all } i \}$$

is ■ equidimensional, ■ connected, ■ components indexed by $\text{SYT}(\eta)$

Combinatorics of Springer Fibers S_η


- Components are indexed by standard Young tableaux of shape η
- Equidimensional of dimension $\sum_i (i-1)\eta_i$

0	0	0
1	1	1
2	2	2

$\leadsto S_{(3,3,3)}$ is 9-dimensional

- $H^*(S_\eta)$ carries an \mathfrak{S}_n -action and
 - gives rise to Hall–Littlewood polynomials
 - top graded piece of $H^*(S_\eta)$ is the Specht module S^η

Geometry of Springer Fibers



2 Column Tableau Descriptions

Let $T \in \text{SYT}(k \times 2)$ with corresponding Springer component F_T . Define:

$$\tau(T) = \{j \in \text{col}_1(T) \mid j + 1 \in \text{col}_2(T)\}$$

Then,

$$F_T \text{ smooth} \stackrel{\text{Fresse, Melnikov ('11)}}{\iff} \begin{array}{l} |\tau(T)| < 3 \text{ or} \\ |\tau(T)| = 3 + \text{another condition} \end{array}$$

Fresse, Melnikov, Sakas-Obeid ('15). If F_T is smooth, it is an *iterated fibre bundle* with base

$$(\text{Fl}(a+b) \times \text{Fl}(b+c), \text{Gr}(a, a+c), \mathbf{P}^b, \dots, \mathbf{P}^{k-1}),$$

where:

τ	a	b	c
$\{\alpha\}$	$\alpha - k$	k	$k - \alpha$
$\{\alpha < \beta\}$	$k - (\beta - \alpha)$	$\beta - k$	$k - \alpha$
$\{\alpha < \beta < \gamma\}$	$k - (\gamma - \beta)$	$(\gamma - \alpha) - k$	$k - (\beta - \alpha)$

2 Column Tableau Examples

1	5
2	7
3	8
4	10
6	11
9	12


singular

1	6
2	7
3	8
4	10
5	11
9	12


smooth

$(\mathrm{Fl}(5) \times \mathrm{Fl}(4), \mathrm{Gr}(2, 3), \mathbf{P}^3, \mathbf{P}^4, \mathbf{P}^5)$


Geometry of Springer Fibers



2 Column Web Examples



cycle \implies singular




acyclic \implies smooth

$(\text{Fl}(5) \times \text{Fl}(4), \text{Gr}(2,3), \mathbf{P}^3, \mathbf{P}^4, \mathbf{P}^5)$

Summary

[C. 26+] Smooth components of two column Springer fibers...

- correspond to acyclic webs
- geometry described by claw sizes
- Poincaré polynomial is rotation/reflection invariant
- $k + 2\binom{k}{3}$ smooth components for $S_{(2^k)}$



arXiv 2602.16910